

Goldschmidt Conference, Paris, France, August 13-18, 2017

Does $p\text{CO}_2$ affect carbon isotope discrimination on evolutionary timescales?

Ying Cui¹ and Brian A. Schubert²

¹*Department of Earth Sciences, Dartmouth College, Hanover, NH 03755*

²*School of Geosciences, University of Louisiana at Lafayette, Lafayette, LA 70504*

Annual tree-ring datasets and growth chamber experiments suggest that the concentration of carbon dioxide in the atmosphere ($p\text{CO}_2$) have an important effect on the carbon isotope fractionation between plant tissue and the atmosphere ($\Delta^{13}\text{C}$). This effect has been identified in Quaternary-aged speleothems and bulk organic matter and within carbon isotope excursion events such as the Paleocene- Eocene Thermal Maximum. Identification of this effect, however, on longer timescales has proven difficult. In order to test whether changes in $p\text{CO}_2$ affect carbon isotope discrimination over evolutionary timescales, we compiled >2000 $\delta^{13}\text{C}$ values measured on plant tissues, bulk organic matter, and specific organic compounds (e.g., *n*-alkanes and *n*-alkanoic acids) spanning the last 65 million years. We derive $\Delta^{13}\text{C}$ using the Cenozoic atmospheric $\delta^{13}\text{C}$ values and our compiled dataset, and show that $\Delta^{13}\text{C}$ follows pattern of $p\text{CO}_2$ based on several other proxies. This is consistent with the fundamental photorespiration effect on $\Delta^{13}\text{C}$ observed in field experiments and growth chambers. Therefore, we reconstruct $p\text{CO}_2$ for the last 65 million years using the new C_3 plant $p\text{CO}_2$ proxy approach and assess the uncertainties using Monte Carlo error propagation. The data suggest elevated $p\text{CO}_2$ during the early Eocene climate optimum (~52-50 Ma) and the Miocene Climate Optimum (~17-15 Ma), and a decline in $p\text{CO}_2$ from early Pliocene into the onset of Pleistocene glacial-interglacial cycles (5-1 Ma). The reconstruction also resolves the $p\text{CO}_2$ oscillation between ~150 and 300 ppmv during the late Pleistocene (1 – 0 Ma), consistent with the ice core $p\text{CO}_2$ records. These results suggest the need to account for the underlying effect of $p\text{CO}_2$ when interpreting $\Delta^{13}\text{C}$ across evolutionary timescales.